Chamorchis

Habitus and growth type

Plant height [m]: **0.09**
Life span: **Perennial**
Life form: **Geophyte**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **0**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **initial or partial mycoheterotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **2.3**
Moisture indicator value: **4.5**
Reaction indicator value: **8.3**
Nutrient indicator value: **2.7**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.55**
Disturbance frequency (herb layer): **1.63**
Disturbance severity: **0.28**
Disturbance severity (herb layer): **0.28**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.01**
Grazing pressure: **0.25**
Soil disturbance: **0.08**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **AB (KOB) *Carici rupestris-Kobresietea bellardii***, **CU (SES) *Elyno-Seslerietea***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Oromediterranean scrub**

Distribution

Continentality: **3**
Continentality amplitude: **3**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz