Neatostema

Habitus and growth type

Plant height [m]: **0.13**
Life span: **Annual**
Life form: **Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **16.07**

Flower

Flowering period: **February-May**

Fruit, seed and dispersal

Seed mass [mg]: **1.23**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **10**
Moisture indicator value: **2**
Reaction indicator value: **7.5**
Nutrient indicator value: **5.5**
Salinity indicator value: **0.5**

Disturbance indicator values

Disturbance frequency: **1.47**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.83**
Disturbance severity: **0.73**
Disturbance severity (herb layer): **0.67**
Mowing frequency: **0.2**
Grazing pressure: **0.4**
Soil disturbance: **0.32**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FJ (TRA) *Stipo-Trachynietea distachyae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R1F Mediterranean annual-rich dry grassland**
Constant species of EUNIS habitats: **R1F Mediterranean annual-rich dry grassland**

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz