Santolina rosmarinifolia aggr.

Habitus and growth type

Plant height [m]: **0.4**
Life span: **Perennial**
Life form: **Chamaephyte**
Spinescence: **not spinescent**

Fruit, seed and dispersal

Seed mass [mg]: **0.3**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **7.5**
Moisture indicator value: **1**
Reaction indicator value: **8**
Nutrient indicator value: **3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.73**
Disturbance frequency (herb layer): **1.36**
Disturbance severity: **0.36**
Disturbance severity (herb layer): **0.22**
Mowing frequency: **0.04**
Grazing pressure: **0.32**
Soil disturbance: **0.12**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **S73 Western Mediterranean mountain hedgehog-heath**

Broad habitat

Occurrence in broad habitats: **Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz