Cuscuta palaestina aggr.

Habitus and growth type

Plant height [m]: **0.42**
Life span: **Annual**
Life form: **Therophyte, Herbaceous liana**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **parasitic vine**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**

Ellenberg-type indicator values

Light indicator value: **7.7**
Temperature indicator value: **9**
Moisture indicator value: **4**
Reaction indicator value: **7**
Nutrient indicator value: **6**
Salinity indicator value: **2.7**

Disturbance indicator values

Disturbance frequency: **0.99**
Disturbance frequency (herb layer): **1.95**
Disturbance severity: **0.78**
Disturbance severity (herb layer): **0.5**
Mowing frequency: **0.17**
Grazing pressure: **0.36**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Soil disturbance: **0.31**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **S63 Eastern garrigue**, **S72 Eastern Mediterranean spiny heath (phrygana)**

Constant species of EUNIS habitats: **S63 Eastern garrigue**

Broad habitat

Occurrence in broad habitats: **Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz