Melampyrum nemorosum aggr.

Habitus and growth type

Plant height [m]: **0.4**
Life span: **Annual**
Life form: **Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **38.18**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **7.94**
Dispersal mode: **Myrmecochory**
Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **euphytoid hemiparasite**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5.1**
Moisture indicator value: **4.6**
Reaction indicator value: **5.4**
Nutrient indicator value: **3.8**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.13**
Disturbance frequency (herb layer): **1.38**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity: **0.65**
Disturbance severity (herb layer): **0.19**
Mowing frequency: **0.09**
Grazing pressure: **0.2**
Soil disturbance: **0.12**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **R51 Thermophilous forest fringe of base-rich soils**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz