

# *Scandix pecten-veneris* aggr.

## Habitus and growth type

Plant height [m]: **0.24**

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **16.68**

## Flower

Flowering period: **March-June**

## Fruit, seed and dispersal

Seed mass [mg]: **13.72**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **1, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.7**

Moisture indicator value: **3.2**

Reaction indicator value: **7.6**

Nutrient indicator value: **4.5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.13**

Disturbance frequency (herb layer): **2.02**

Disturbance severity: **0.81**

Disturbance severity (herb layer): **0.72**

Mowing frequency: **0.38**

Grazing pressure: **0.22**

Soil disturbance: **0.56**

## **Habitat and sociology**

EUNIS habitat

Diagnostic species of EUNIS habitats: [S63 Eastern garrigue](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline),  
Oromediterranean scrub, Scrub, Sparsely vegetated (incl. rock and scree),  
Synanthropic**