

# ***Viola tricolor* aggr.**

## **Habitus and growth type**

Plant height [m]: **0.25**

Life span: **Annual, Biennial or short-lived, Perennial**

Life form: **Hemicryptophyte, Therophyte**

## **Leaf**

Specific leaf area [mm<sup>2</sup>/mg]: **27.81**

## **Flower**

Flowering period: **April-July**

## **Fruit, seed and dispersal**

Seed mass [mg]: **0.71**

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

### Environmental relationships

Substrate reaction relationship: **Slightly acidic to near-neutral**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.8**

Reaction indicator value: **5.1**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.2**

Disturbance frequency (herb layer): **1.74**

Disturbance severity: **0.54**

Disturbance severity (herb layer): **0.51**

Mowing frequency: **0.45**

Grazing pressure: **0.26**

Soil disturbance: **0.41**

## **Habitat and sociology**

EUNIS habitat

Diagnostic species of EUNIS habitats: [S34 Balkan-Anatolian submontane genistoid scrub](#)

Constant species of EUNIS habitats: [S34 Balkan-Anatolian submontane genistoid scrub](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**