# *Achillea pyrenaica*

## Habitus and growth type

Plant height [m]: **0.76**
Life span: **Perennial**
Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [$mm^2$/mg]: **16.48**

## Fruit, seed and dispersal

Seed mass [mg]: **0.3**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **3**
Moisture indicator value: **6.5**
Reaction indicator value: **3**
Nutrient indicator value: **3**
Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.6**
Disturbance frequency (herb layer): **1.88**
Disturbance severity: **0.51**
Disturbance severity (herb layer): **0.5**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Mowing frequency: **1.04**
Grazing pressure: **0.24**
Soil disturbance: **0.22**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **CM (MOL) *Molinio-Arrhenatheretea*, CS (MUL) *Mulgedio-Aconitetea***

### EUNIS habitat

Diagnostic species of EUNIS habitats: **R35 Moist or wet mesotrophic to eutrophic hay meadow, R37 Temperate and boreal moist or wet oligotrophic grassland**

### Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland**

## Distribution

Continentality amplitude: **5**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*