# *Allium pallens*

### Habitus and growth type

Plant height [m]: **0.53**
Life span: **Perennial**
Life form: **Geophyte**

### Flower

Flowering period: **May-July**

### Fruit, seed and dispersal

Seed mass [mg]: **1.62**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### Taxon origin

Origin in Europe: **native**

### Ecology

#### Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

#### Ellenberg-type indicator values

Light indicator value: **7**
Temperature indicator value: **9.3**
Moisture indicator value: **3**
Reaction indicator value: **7**
Nutrient indicator value: **5**
Salinity indicator value: **0**

#### Disturbance indicator values

Disturbance frequency: **0.69**
Disturbance frequency (herb layer): **1.53**
Disturbance severity: **0.63**
Disturbance severity (herb layer): **0.51**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Mowing frequency: **0.35**
Grazing pressure: **0.38**
Soil disturbance: **0.19**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **FF (LYG) *Lygeo sparti-Stipetea tenacissimae***

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*