

Allium subhirsutum

Habitus and growth type

Plant height [m]: **0.29**

Life span: **Perennial**

Life form: **Geophyte**

Flower

Flowering period: **February-May**

Fruit, seed and dispersal

Seed mass [mg]: **1.45**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **9.3**

Moisture indicator value: **4**

Reaction indicator value: **7.3**

Nutrient indicator value: **6.3**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **0.2**

Disturbance frequency (herb layer): **1.36**

Disturbance severity: **0.7**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0.06**

Grazing pressure: **0.27**

Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CB \(PUB\) Quercetea pubescentis](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**