# *Ambrosia trifida*

### Habitus and growth type

Plant height [m]: **1.3**
Life span: **Annual**
Life form: **Therophyte**

### Leaf

Specific leaf area [$mm^2$/mg]: **24.11**

### Flower

Flowering period: **June-September**

### Fruit, seed and dispersal

Seed mass [mg]: **14.06**
Dispersal mode: **Anemochory, Anthropochory**
Dispersal distance class: **5, 7**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### Taxon origin

Origin in Europe: **neophyte**

### Ecology

#### Disturbance indicator values

Disturbance frequency: **1.93**
Disturbance frequency (herb layer): **1.93**
Disturbance severity: **0.86**
Disturbance severity (herb layer): **0.86**
Mowing frequency: **1.17**
Grazing pressure: **0.11**
Soil disturbance: **0.76**

### Habitat and sociology

#### Syntaxon

Diagnostic species of phytosociological classes: **QB (SIS) *Sisymbrietea***

#### Broad habitat

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Occurrence in broad habitats: **Wetland, Sparsely vegetated (incl. rock and scree), Synanthropic**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*