# *Anacyclus radiatus*

## Habitus and growth type

Plant height [m]: **0.38**
Life span: **Annual**
Life form: **Therophyte**

## Flower

Flowering period: **April-August**

## Fruit, seed and dispersal

Seed mass [mg]: **1.07**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **8.5**
Moisture indicator value: **3.5**
Reaction indicator value: **6**
Nutrient indicator value: **5.5**
Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.99**
Disturbance frequency (herb layer): **2.21**
Disturbance severity: **0.85**
Disturbance severity (herb layer): **0.85**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Mowing frequency: **0.74**
Grazing pressure: **0.17**
Soil disturbance: **0.53**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **QC (CHE) *Chenopodietea***

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Synanthropic**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*