

# *Anthemis secundiramea*

## Habitus and growth type

Plant height [m]: **0.16**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **March-June**

## Fruit, seed and dispersal

Seed mass [mg]: **0.15**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

### Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **9.3**

Moisture indicator value: **3.8**

Reaction indicator value: **6.8**

Nutrient indicator value: **6**

Salinity indicator value: **2**

### Disturbance indicator values

Disturbance frequency: **0.9**

Disturbance frequency (herb layer): **0.93**

Disturbance severity: **0.51**

Disturbance severity (herb layer): **0.5**

Mowing frequency: **0.11**

Grazing pressure: **0.27**

Soil disturbance: **0.23**

### **Habitat and sociology**

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Coastal cliff, Scrub, Sparsely vegetated (incl. rock and scree)**