# *Anthriscus caucalis*

### Habitus and growth type

Plant height [m]: **0.45**
Life span: **Annual**
Life form: **Therophyte**

### Leaf

Specific leaf area [$mm^2$/mg]: **32.98**

### Flower

Flowering period: **April-July**

### Fruit, seed and dispersal

Seed mass [mg]: **1.62**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### Taxon origin

Origin in Europe: **native**

### Ecology

#### Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

#### Ellenberg-type indicator values

Light indicator value: **7.3**
Reaction indicator value: **6**
Nutrient indicator value: **6.5**
Salinity indicator value: **0.1**

#### Disturbance indicator values

Disturbance frequency: **0.72**
Disturbance frequency (herb layer): **1.49**
Disturbance severity: **0.63**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Disturbance severity (herb layer): **0.5**
Mowing frequency: **0.29**
Grazing pressure: **0.21**
Soil disturbance: **0.35**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **QB (SIS) *Sisymbrietea*, QG (EPI) *Epilobietea angustifolii***

### Broad habitat

Occurrence in broad habitats: **Forest, Synanthropic**

## Distribution

Continentality: **3**
Continentality amplitude: **3**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*