Anthriscus sylvestris

Habitus and growth type

Plant height [m]: **0.96**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **33.52**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **3.88**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Reaction indicator value: **6.6**
Nutrient indicator value: **7.4**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.69**
Disturbance frequency (herb layer): **1.55**
Disturbance severity: **0.57**
Disturbance severity (herb layer): **0.45**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.53**
Grazing pressure: **0.16**
Soil disturbance: **0.27**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **R22 Low and medium altitude hay meadow**, **V39 Mesic perennial anthropogenic herbaceous vegetation**
Constant species of EUNIS habitats: **N1D Atlantic and Baltic broad-leaved coastal dune forest**, **R22 Low and medium altitude hay meadow**, **R55 Lowland moist or wet tall-herb and fern fringe**, **V39 Mesic perennial anthropogenic herbaceous vegetation**

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **6**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz