Artemisia campestris

Habitus and growth type

Plant height [m]: **0.61**
Life span: **Perennial**
Life form: **Chamaephyte, Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **16.31**

Flower

Flowering period: **July-October**

Fruit, seed and dispersal

Seed mass [mg]: **0.16**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**

Ellenberg-type indicator values

Light indicator value: **8.6**
Moisture indicator value: **3.1**
Reaction indicator value: **5.7**
Salinity indicator value: **0.8**

Disturbance indicator values

Disturbance frequency: **1.34**
Disturbance frequency (herb layer): **1.79**
Disturbance severity: **0.48**
Disturbance severity (herb layer): **0.45**
Mowing frequency: **0.47**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.29**
Soil disturbance: **0.28**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **R11 Pannonian and Pontic sandy steppe**, **R1P Oceanic to subcontinental inland sand grassland on dry acid and neutral soils**, **U72 Unvegetated or sparsely vegetated gravel bar in the Mediterranean region**
Constant species of EUNIS habitats: **N15 Atlantic and Baltic coastal dune grassland (grey dune)**, **N17 Black Sea coastal dune grassland (grey dune)**, **R11 Pannonian and Pontic sandy steppe**, **R13 Cryptogam- and annual-dominated vegetation on calcareous and ultramafic rock outcrops**, **R16 Perennial rocky grassland of Central and South-Eastern Europe**, **R1B Continental dry grassland (true steppe)**, **R1P Oceanic to subcontinental inland sand grassland on dry acid and neutral soils**, **S36 Low steppic scrub**, **U33 Temperate, lowland to montane siliceous inland cliff**, **U72 Unvegetated or sparsely vegetated gravel bar in the Mediterranean region**

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Coastal cliff, Grassland (non-alpine, non-saline), Semi-desert, Scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz