

# *Artemisia taurica*

## Habitus and growth type

Life span: **Perennial**

Life form: **Chamaephyte, Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **5.5**

Moisture indicator value: **4**

Disturbance indicator values

Disturbance frequency: **1.87**

Disturbance frequency (herb layer): **1.87**

Disturbance severity: **0.51**

Disturbance severity (herb layer): **0.5**

Mowing frequency: **0.33**

Grazing pressure: **0.41**

Soil disturbance: **0.26**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1C Desert steppe](#), [S67 Aralo-Caspian semi-desert](#)

Constant species of EUNIS habitats: [S67 Aralo-Caspian semi-desert](#)

## Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Semi-desert**