

# *Arum pictum*

## Habitus and growth type

Plant height [m]: **0.4**

Life span: **Perennial**

Life form: **Geophyte**

## Flower

Flowering period: **September-November**

## Fruit, seed and dispersal

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **6.5**

Temperature indicator value: **8**

Moisture indicator value: **4.5**

Reaction indicator value: **6.5**

Nutrient indicator value: **4.5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.25**

Disturbance frequency (herb layer): **1.44**

Disturbance severity: **0.67**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0.03**

Grazing pressure: **0.27**

Soil disturbance: **0.16**

### **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [FA \(QUI\) Quercetea ilicis](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Forest, Sparsely vegetated (incl. rock and scree)**