Bombycilaena discolor

Habitus and growth type

Plant height [m]: **0.14**
Life span: **Annual**
Life form: **Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **18.14**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values
Temperature indicator value: **9**

Disturbance indicator values
Disturbance frequency: **1.25**
Disturbance frequency (herb layer): **2.12**
Disturbance severity: **0.51**
Disturbance severity (herb layer): **0.5**
Mowing frequency: **0.21**
Grazing pressure: **0.31**
Soil disturbance: **0.24**

Habitat and sociology

Syntaxon
Diagnostic species of phytosociological classes: **CJ (COR) *Koelerio-Corynephoretea***

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

canescentis*, FI (TUB) *Helianthemetea guttati*, FJ (TRA) *Stipo-Trachynietea distachyae

EUNIS habitat

Diagnostic species of EUNIS habitats: **S65 Mediterranean gypsum scrub**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz