

Bombycilaena discolor

Habitus and growth type

Plant height [m]: **0.14**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **18.14**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Temperature indicator value: **9**

Disturbance indicator values

Disturbance frequency: **1.25**

Disturbance frequency (herb layer): **2.12**

Disturbance severity: **0.51**

Disturbance severity (herb layer): **0.5**

Mowing frequency: **0.21**

Grazing pressure: **0.31**

Soil disturbance: **0.24**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CJ \(COR\) Koelerio-Coryneporetea](#)

canescentis, FI (TUB) *Helianthemetea guttati*, FJ (TRA) *Stipo-Trachynietea distachyae*

EUNIS habitat

Diagnostic species of EUNIS habitats: [S65 Mediterranean gypsum scrub](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**