

Brassica tournefortii

Habitus and growth type

Plant height [m]: **0.4**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **10.61**

Flower

Flowering period: **January-May**

Fruit, seed and dispersal

Seed mass [mg]: **1.4**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.7**

Temperature indicator value: **10.7**

Moisture indicator value: **3.3**

Reaction indicator value: **8**

Nutrient indicator value: **5.7**

Salinity indicator value: **1**

Disturbance indicator values

Disturbance frequency: **1.11**

Disturbance frequency (herb layer): **1.94**

Disturbance severity: **0.57**

Disturbance severity (herb layer): **0.47**

Mowing frequency: **0.13**

Grazing pressure: **0.22**

Soil disturbance: **0.42**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FJ \(TRA\) *Stipo-Trachynietea distachyae*](#), [QC \(CHE\) *Chenopodietea*](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Synanthropic**