Bromus arvensis

Habitus and growth type

Plant height [m]: **0.46**
Life span: **Annual**
Life form: **Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **19.2**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **2.31**
Dispersal mode: **Local non-specific dispersal, Anthropochory**
Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**
Reaction indicator value: **7.9**
Nutrient indicator value: **5.2**
Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **1.24**
Disturbance frequency (herb layer): **1.86**
Disturbance severity: **0.68**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity (herb layer): **0.64**
Mowing frequency: **0.68**
Grazing pressure: **0.22**
Soil disturbance: **0.43**

Habitat and sociology

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Synanthropic**

Distribution

Continentality amplitude: **6**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz