Bupleurum fruticescens

Habitus and growth type

Plant height [m]:
Life span: **Perennial**
Life form: **Chamaephyte**
Spinescence: **not spinescent**

Leaf

Specific leaf area [mm^2/mg]: **8.25**

Fruit, seed and dispersal

Seed mass [mg]: **2.42**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.3**
Temperature indicator value: **8.2**
Moisture indicator value: **2.5**
Reaction indicator value: **7.1**
Nutrient indicator value: **2.8**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.36**
Disturbance frequency (herb layer): **1.42**
Disturbance severity: **0.65**
Disturbance severity (herb layer): **0.21**
Mowing frequency: **0.04**
Grazing pressure: **0.36**
Soil disturbance: **0.14**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **S61 Western basiphilous garrigue**, **S73 Western Mediterranean mountain hedgehog-heath**

Constant species of EUNIS habitats: **S61 Western basiphilous garrigue**, **S73 Western Mediterranean mountain hedgehog-heath**

Dominant species of EUNIS habitats: **S73 Western Mediterranean mountain hedgehog-heath**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Oromediterranean scrub, Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz