Bupleurum longifolium

Habitus and growth type

Plant height [m]: **0.64**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **June-July**

Fruit, seed and dispersal

Seed mass [mg]: **2.92**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5.4**
Moisture indicator value: **4.5**
Reaction indicator value: **8.5**
Nutrient indicator value: **5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.29**
Disturbance frequency (herb layer): **1.41**
Disturbance severity: **0.58**
Disturbance severity (herb layer): **0.24**
Mowing frequency: **0.12**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.19**
Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **BC (BRA) *Brachypodio pinnati-Betuletea pendulae***, **CL (GER) *Trifolio-Geranietea sanguinei***, **CU (SES) *Elyno-Seslerietea***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub, Forest**

Distribution

Continentality: **4**
Continentality amplitude: **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz