# *Camelina microcarpa*

## Habitus and growth type

Plant height [m]: **0.35**
Life span: **Annual**
Life form: **Therophyte**

## Leaf

Specific leaf area [$mm^2$/mg]: **19.99**

## Flower

Flowering period: **April-June**

## Fruit, seed and dispersal

Seed mass [mg]: **0.31**
Dispersal mode: **Local non-specific dispersal, Anthropochory**
Dispersal distance class: **2, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.4**
Moisture indicator value: **3.9**
Reaction indicator value: **7.3**
Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.64**
Disturbance frequency (herb layer): **1.94**
Disturbance severity: **0.69**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Disturbance severity (herb layer): **0.68**
Mowing frequency: **0.59**
Grazing pressure: **0.24**
Soil disturbance: **0.5**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea*, QA (PAR) *Papaveretea rhoeadis***

### EUNIS habitat

Diagnostic species of EUNIS habitats: **S67 Aralo-Caspian semi-desert**

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Synanthropic**

## Distribution

Continentality: **7**
Continentality amplitude: **5**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*