

Campanula dichotoma

Habitus and growth type

Plant height [m]: **0.12**

Life span: **Annual**

Life form: **Therophyte**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **8**

Moisture indicator value: **4**

Reaction indicator value: **5**

Nutrient indicator value: **5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.45**

Disturbance frequency (herb layer): **1.67**

Disturbance severity: **0.82**

Disturbance severity (herb layer): **0.77**

Mowing frequency: **0.51**

Grazing pressure: **0.21**

Soil disturbance: **0.43**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FI (TUB) *Helianthemetea guttati***

Broad habitat

Occurrence in broad habitats: **Spring, Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**