

# *Campanula morettiana*

## Habitus and growth type

Plant height [m]: **0.05**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **July-August**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **4**

Temperature indicator value: **2.5**

Moisture indicator value: **4**

Reaction indicator value: **7**

Nutrient indicator value: **1**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.55**

Disturbance frequency (herb layer): **0.55**

Disturbance severity: **0.11**

Disturbance severity (herb layer): **0.11**

Mowing frequency: **0**

Grazing pressure: **0.02**

Soil disturbance: **0.01**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [KC \(ASP\) \*Asplenietea trichomanis\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [U36 Temperate high-mountain base-rich inland cliff](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Sparsely vegetated (incl. rock and scree)**