

Campanula rotundifolia

Habitus and growth type

Plant height [m]: **0.3**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **25.45**

Flower

Flowering period: **April-October**

Fruit, seed and dispersal

Seed mass [mg]: **0.06**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Moisture indicator value: **4**

Nutrient indicator value: **2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.69**

Disturbance frequency (herb layer): **1.53**

Disturbance severity: **0.5**

Disturbance severity (herb layer): **0.37**

Mowing frequency: **0.37**

Grazing pressure: **0.25**

Soil disturbance: **0.2**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CL \(GER\) *Trifolio-Geranietea sanguinei*](#), [DA \(FES\) *Festuco-Brometea*](#), [FP \(ONO\) *Festuco hystricis-Ononidetea striatae*](#), [FR \(DAP\) *Daphno-Festucetea*](#), [KC \(ASP\) *Asplenetetea trichomanis*](#), [KE \(THL\) *Thlaspietea rotundifolii*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1S Heavy-metal grassland in Western and Central Europe](#), [U21 Boreal and Arctic siliceous scree and block field](#), [U33 Temperate, lowland to montane siliceous inland cliff](#), [U35 Boreal and Arctic base-rich inland cliff](#)

Constant species of EUNIS habitats: [R16 Perennial rocky grassland of Central and South-Eastern Europe](#), [R1A Semi-dry perennial calcareous grassland \(meadow steppe\)](#), [R1M Lowland to montane, dry to mesic grassland usually dominated by *Nardus stricta*](#), [R1S Heavy-metal grassland in Western and Central Europe](#), [R23 Mountain hay meadow](#), [R42 Boreal and Arctic acidophilous alpine grassland](#), [T34 Temperate subalpine *Larix*, *Pinus cembra* and *Pinus uncinata* forest](#), [T36 Temperate and submediterranean montane *Pinus sylvestris*-*Pinus nigra* forest](#), [T3H *Larix* light taiga](#), [U21 Boreal and Arctic siliceous scree and block field](#), [U23 Temperate, lowland to montane siliceous scree](#), [U25 Boreal and Arctic base-rich scree and block field](#), [U32 Temperate high-mountain siliceous inland cliff](#), [U33 Temperate, lowland to montane siliceous inland cliff](#), [U34 Mediterranean siliceous inland cliff](#), [U35 Boreal and Arctic base-rich inland cliff](#), [U37 Temperate, lowland to montane base-rich inland cliff](#), [U38 Mediterranean base-rich inland cliff](#), [U3A Temperate ultramafic inland cliff](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Spring, Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Heathland, Oromediterranean scrub, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality amplitude: 7