Capparis orientalis

Habitus and growth type

Life span: **Perennial**
Life form: **Phanerophyte, Shrub**

Fruit, seed and dispersal

Dispersal mode: **Endozoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **7.5**
Temperature indicator value: **9.3**
Moisture indicator value: **2**
Reaction indicator value: **8**
Nutrient indicator value: **5.5**
Salinity indicator value: **1.5**

Disturbance indicator values

Disturbance frequency: **0.63**
Disturbance frequency (herb layer): **0.64**
Disturbance severity: **0.27**
Disturbance severity (herb layer): **0.25**
Mowing frequency: **0.1**
Grazing pressure: **0.1**
Soil disturbance: **0.07**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **KD (CYM) *Cymbalario-Parietarietea diffusae***

Broad habitat

Occurrence in broad habitats: **Synanthropic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz