

Cardamine resedifolia

Habitus and growth type

Plant height [m]: **0.07**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **20.3**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.18**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **1.8**

Moisture indicator value: **5**

Reaction indicator value: **3**

Nutrient indicator value: **2.5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.02**

Disturbance frequency (herb layer): **1.37**
Disturbance severity: **0.31**
Disturbance severity (herb layer): **0.28**
Mowing frequency: **0.01**
Grazing pressure: **0.18**
Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [KC \(ASP\) *Asplenietea trichomanis*](#),
[KE \(THL\) *Thlaspietea rotundifolii*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [U22 Temperate high-mountain siliceous scree](#), [U24 Mediterranean siliceous scree](#), [U32 Temperate high-mountain siliceous inland cliff](#)

Constant species of EUNIS habitats: [U22 Temperate high-mountain siliceous scree](#), [U24 Mediterranean siliceous scree](#), [U32 Temperate high-mountain siliceous inland cliff](#)

Broad habitat

Occurrence in broad habitats: **Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **4**
Continentality amplitude: **2**