Carex globularis

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte, Geophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Ellenberg-type indicator values

Light indicator value: **7**
Temperature indicator value: **3.8**
Moisture indicator value: **7.6**

Disturbance indicator values

Disturbance frequency: **0.14**
Disturbance frequency (herb layer): **1.34**
Disturbance severity: **0.64**
Disturbance severity (herb layer): **0.11**
Mowing frequency: **0.01**
Grazing pressure: **0.18**
Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **BA (PIC) *Vaccinio-Piceetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T3J Pinus and Larix mire forest**, **T3K Picea mire forest**
Constant species of EUNIS habitats: **T3J Pinus and Larix mire forest**, **T3K Picea mire**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

forest

Broad habitat

Occurrence in broad habitats: **Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz