Carex riparia

Habitus and growth type

Plant height [m]: **1.03**
Life span: **Perennial**
Life form: **Hemicryptophyte, Geophyte**

Leaf

Specific leaf area [mm^2/mg]: **11.59**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **2.29**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.9**
Temperature indicator value: **5.8**
Moisture indicator value: **9.4**
Reaction indicator value: **6.7**
Nutrient indicator value: **5.4**
Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **0.8**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.59**
Disturbance severity: **0.37**
Disturbance severity (herb layer): **0.29**
Mowing frequency: **0.44**
Grazing pressure: **0.15**
Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **OD (PHR) *Phragmito-Magnocaricetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **Q53 Tall-sedge bed**
Constant species of EUNIS habitats: **Q53 Tall-sedge bed**
Dominant species of EUNIS habitats: **Q53 Tall-sedge bed**

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Grassland (non-alpine, non-saline), Scrub, Forest**

Distribution

Continentality amplitude: **9**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz