

Centaureum pulchellum

Habitus and growth type

Plant height [m]: **0.1**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **47.01**

Flower

Flowering period: **April-October**

Fruit, seed and dispersal

Seed mass [mg]: **0.01**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8.7**

Moisture indicator value: **6.7**

Reaction indicator value: **7.9**

Nutrient indicator value: **3.4**

Salinity indicator value: **1.1**

Disturbance indicator values

Disturbance frequency: **1.53**

Disturbance frequency (herb layer): **1.97**

Disturbance severity: **0.56**

Disturbance severity (herb layer): **0.54**

Mowing frequency: **0.3**

Grazing pressure: **0.42**

Soil disturbance: **0.23**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [OC \(ISO\) Isoëto-Nanojuncetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [MA222 Atlantic upper saltmarsh](#), [R32 Mediterranean short moist grassland of lowlands](#)

Constant species of EUNIS habitats: [MA222 Atlantic upper saltmarsh](#), [R32 Mediterranean short moist grassland of lowlands](#)

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Wetland, Grassland (non-alpine, non-saline), Saline vegetation, Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **10**