

# *Cerintho minor*

## Habitus and growth type

Plant height [m]: **0.37**

Life span: **Annual, Biennial or short-lived, Perennial**

Life form: **Hemicryptophyte, Therophyte**

## Flower

Flowering period: **April-August**

## Fruit, seed and dispersal

Seed mass [mg]: **9.06**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.6**

Moisture indicator value: **4.3**

Reaction indicator value: **6.8**

Nutrient indicator value: **4.2**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.01**

Disturbance frequency (herb layer): **1.55**

Disturbance severity: **0.59**

Disturbance severity (herb layer): **0.52**

Mowing frequency: **0.55**

Grazing pressure: **0.22**

Soil disturbance: **0.29**

### **Habitat and sociology**

Broad habitat

Occurrence in broad habitats: **Spring, Wetland, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

### **Distribution**

Continentality: 7

Continentality amplitude: 3