Corylus avellana

Habitus and growth type

Plant height [m]: **6.21**
Life span: **Perennial**
Life form: **Phanerophyte, Shrub**

Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**
Leaf phenology (woody plants only): **Winter deciduous**
Specific leaf area [mm^2/mg]: **23.63**

Flower

Flowering period: **February-April**

Fruit, seed and dispersal

Seed mass [mg]: **984.14**
Dispersal mode: **Dyszoochory, Anthropochory**
Dispersal distance class: **6, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Moisture indicator value: **5.6**
Reaction indicator value: **5.9**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.07**
Disturbance frequency (herb layer): **1.34**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.16**
Mowing frequency: **0.01**
Grazing pressure: **0.19**
Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CD (RHA) *Crataego-Prunetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S37 Corylus avellana scrub**, **S38 Temperate forest clearing scrub**, **T13 Temperate hardwood riparian forest**, **T1D Southern European mountain Betula and Populus tremula forest on mineral soils**, **T1E Carpinus and Quercus mesic deciduous forest**, **T1F Ravine forest**, **T32 Temperate mountain Abies forest**, **T3C Taxus baccata forest**

Constant species of EUNIS habitats: **S35 Temperate and submediterranean thorn scrub**, **S37 Corylus avellana scrub**, **S38 Temperate forest clearing scrub**, **S52 Submediterranean pseudomaquis**, **T12 Alnus glutinosa-Alnus incana forest on riparian and mineral soils**, **T13 Temperate hardwood riparian forest**, **T17 Fagus forest on non-acid soils**, **T19 Temperate and submediterranean thermophilous deciduous forest**, **T1B Acidophilous Quercus forest**, **T1D Southern European mountain Betula and Populus tremula forest on mineral soils**, **T1E Carpinus and Quercus mesic deciduous forest**, **T1F Ravine forest**, **T1H Broadleaved deciduous plantation of non site-native trees**, **T22 Mainland laurophyllous forest**, **T27 Ilex aquifolium forest**, **T31 Temperate mountain Picea forest**, **T32 Temperate mountain Abies forest**, **T36 Temperate and submediterranean montane Pinus sylvestris-Pinus nigra forest**, **T3C Taxus baccata forest**, **T3F Dark taiga**, **T3M Coniferous plantation of non site-native trees**

Dominant species of EUNIS habitats: **S37 Corylus avellana scrub**, **S38 Temperate forest clearing scrub**, **T13 Temperate hardwood riparian forest**, **T1D Southern European mountain Betula and Populus tremula forest on mineral soils**, **T1E Carpinus and Quercus mesic deciduous forest**, **T1F Ravine forest**, **T3C Taxus baccata forest**

Broad habitat

Occurrence in broad habitats: **Scrub, Forest**

Distribution

Continentality amplitude: **5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz