

# *Crassula lycopodioides*

## Habitus and growth type

Plant height [m]: **0.17**

Life span: **Perennial**

Life form: **Chamaephyte**

## Flower

Flowering period: **January-February, September-December**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **1, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Neophyte**

## Ecology

Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **8.6**

Moisture indicator value: **3.1**

Reaction indicator value: **4**

Nutrient indicator value: **2.3**

Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **1.99**

Disturbance frequency (herb layer): **1.99**

Disturbance severity: **0.49**

Disturbance severity (herb layer): **0.49**

Mowing frequency: **0.42**

Grazing pressure: **0.39**

Soil disturbance: **0.33**

## Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **[R1R Mediterranean to Atlantic open, dry, acid and neutral grassland](#)**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**