

Crepis pyrenaica

Habitus and growth type

Plant height [m]: **0.43**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **45**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **3.18**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.5**

Temperature indicator value: **2.8**

Moisture indicator value: **5.5**

Reaction indicator value: **6.3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.46**

Disturbance frequency (herb layer): **1.46**

Disturbance severity: **0.53**

Disturbance severity (herb layer): **0.31**

Mowing frequency: **0.21**

Grazing pressure: **0.19**

Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) *Molinio-Arrhenatheretea*](#),
[CS \(MUL\) *Mulgedio-Aconitetea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R23 Mountain hay meadow](#)

Constant species of EUNIS habitats: [R23 Mountain hay meadow](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub, Forest**

Distribution

Continentality: **4**

Continentality amplitude: **2**