

# *Crepis sibirica*

## Habitus and growth type

Plant height [m]: **0.9**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **3**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **6.3**

Temperature indicator value: **4**

Moisture indicator value: **5.6**

Reaction indicator value: **7**

Nutrient indicator value: **6**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.17**

Disturbance frequency (herb layer): **1.39**

Disturbance severity: **0.64**

Disturbance severity (herb layer): **0.23**

Mowing frequency: **0.1**

Grazing pressure: **0.19**

Soil disturbance: **0.11**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [BB \(ASA\) \*Asaro europaei-Abietetea sibiricae\*](#), [CM \(MOL\) \*Molinio-Arrhenatheretea\*](#), [CS \(MUL\) \*Mulgedio-Aconitetea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [T3H \*Larix light taiga\*](#)

Constant species of EUNIS habitats: [T3H \*Larix light taiga\*](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub, Forest**