

Crepis tectorum

Habitus and growth type

Plant height [m]: **0.44**

Life span: **Annual, Biennial or short-lived**

Life form: **Hemicryptophyte, Therophyte**

Leaf

Specific leaf area [mm²/mg]: **17.2**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.27**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry, Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.8**

Temperature indicator value: **5.8**

Moisture indicator value: **3.9**

Reaction indicator value: **5.5**

Nutrient indicator value: **5.8**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.38**

Disturbance frequency (herb layer): **1.97**
Disturbance severity: **0.7**
Disturbance severity (herb layer): **0.68**
Mowing frequency: **0.5**
Grazing pressure: **0.19**
Soil disturbance: **0.53**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QB \(SIS\) Sisymbrietea](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality: 7
Continentality amplitude: 7