

Dipcadi serotinum

Habitus and growth type

Plant height [m]: **0.25**

Life span: **Perennial**

Life form: **Geophyte**

Spinescence: **not spinescent**

Leaf

Specific leaf area [mm²/mg]: **10.83**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **1.76**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **8**

Moisture indicator value: **3.5**

Reaction indicator value: **7.5**

Nutrient indicator value: **2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.77**
Disturbance frequency (herb layer): **1.6**
Disturbance severity: **0.68**
Disturbance severity (herb layer): **0.54**
Mowing frequency: **0.19**
Grazing pressure: **0.41**
Soil disturbance: **0.22**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FF \(LYG\) *Lygeo sparti-Stipetea tenacissimae*](#), [FI \(TUB\) *Helianthemetea guttati*](#), [GA \(KLE\) *Kleinio neriifoliae-Euphorbieteae canariensis*](#), [GE \(CAN\) *Cytiso-Pinetea canariensis*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**