Diplotaxis virgata

Habitus and growth type

Life span: **Annual**
Life form: **Therophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.1**
Temperature indicator value: **8.3**
Moisture indicator value: **3**
Reaction indicator value: **6.7**
Nutrient indicator value: **5.1**
Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **1.75**
Disturbance frequency (herb layer): **2.07**
Disturbance severity: **0.78**
Disturbance severity (herb layer): **0.77**
Mowing frequency: **0.54**
Grazing pressure: **0.24**
Soil disturbance: **0.45**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QC (CHE) *Chenopodietea***

Broad habitat

Occurrence in broad habitats: **Synanthropic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz