Drimia pancration

Habitus and growth type

Plant height [m]:
Life span: **Perennial**
Life form: **Geophyte**

Flower

Flowering period: **August-October**

Fruit, seed and dispersal

Seed mass [mg]: **2.03**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Oligotrophic**

Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **10**
Moisture indicator value: **1**
Reaction indicator value: **4**
Nutrient indicator value: **2**

Disturbance indicator values

Disturbance frequency: **0.61**
Disturbance frequency (herb layer): **1.41**
Disturbance severity: **0.71**
Disturbance severity (herb layer): **0.45**
Mowing frequency: **0.14**
Grazing pressure: **0.33**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Soil disturbance: **0.19**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **S54 Thermomediterranean arid scrub**

Constant species of EUNIS habitats: **S54 Thermomediterranean arid scrub**

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz