# *Edraianthus tenuifolius*

## Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Ellenberg-type indicator values

Light indicator value: **8.3**
Temperature indicator value: **7.1**
Moisture indicator value: **2.7**
Reaction indicator value: **7.4**
Nutrient indicator value: **2**
Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.73**
Disturbance frequency (herb layer): **1.75**
Disturbance severity: **0.54**
Disturbance severity (herb layer): **0.38**
Mowing frequency: **0.42**
Grazing pressure: **0.42**
Soil disturbance: **0.17**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **CU (SES) *Elyno-Seslerietea***

### EUNIS habitat

Diagnostic species of EUNIS habitats: **R19 Dry steppic submediterranean pasture of the Amphi-Adriatic region**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Constant species of EUNIS habitats: **R19 Dry steppic submediterranean pasture of the Amphi-Adriatic region**

## Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*