Equisetum ramosissimum

Habitus and growth type

Plant height [m]: **0.82**
Life span: **Perennial**
Life form: **Hemicryptophyte, Geophyte**

Flower

Flowering period: **January-October**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate reaction relationship: **Alkaline**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.8**
Reaction indicator value: **7.3**
Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **0.76**
Disturbance frequency (herb layer): **1.6**
Disturbance severity: **0.57**
Disturbance severity (herb layer): **0.46**
Mowing frequency: **0.36**
Grazing pressure: **0.22**
Soil disturbance: **0.23**

Habitat and sociology

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Syntaxon

Diagnostic species of phytosociological classes: **CJ (COR) *Koelerio-Corynephoretea canescentis*, KE (THL) *Thlaspietea rotundifolii***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R31 Mediterranean tall humid inland grassland, T14 Mediterranean and Macaronesian riparian forest, T25 Phoenix theophrasti vegetation**

Constant species of EUNIS habitats: **R31 Mediterranean tall humid inland grassland, T14 Mediterranean and Macaronesian riparian forest, T25 Phoenix theophrasti vegetation**

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality: **7**

Continentality amplitude: **9**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz