Eragrostis barrelieri

Habitus and growth type

Plant height [m]: **0.3**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **May-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.04**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **8.3**
Moisture indicator value: **3.5**
Reaction indicator value: **6.5**
Salinity indicator value: **0**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QD (DIG) *Digitario sanguinalis-Eragrostietea minoris***

Broad habitat

Occurrence in broad habitats: **Sparsely vegetated (incl. rock and scree),**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Synanthropic

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz