

Erodium ciconium

Habitus and growth type

Plant height [m]: **0.42**

Life span: **Annual, Biennial or short-lived**

Life form: **Hemicryptophyte, Therophyte**

Leaf

Specific leaf area [mm²/mg]: **16.73**

Flower

Flowering period: **February-May**

Fruit, seed and dispersal

Seed mass [mg]: **7.98**

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.4**

Moisture indicator value: **2.7**

Reaction indicator value: **7.1**

Nutrient indicator value: **5.4**

Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **1.41**

Disturbance frequency (herb layer): **2.14**

Disturbance severity: **0.78**

Disturbance severity (herb layer): **0.75**

Mowing frequency: **0.65**

Grazing pressure: **0.25**

Soil disturbance: **0.47**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QB \(SIS\) *Sisymbrietea*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Synanthropic**

Distribution

Continentality amplitude: 7