

Euphrasia stricta

Habitus and growth type

Plant height [m]: **0.24**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **22.85**

Flower

Flowering period: **April-October**

Fruit, seed and dispersal

Seed mass [mg]: **0.15**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **euphytoid hemiparasite**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.1**

Temperature indicator value: **5**

Reaction indicator value: **5.4**

Nutrient indicator value: **2.1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.56**

Disturbance frequency (herb layer): **1.94**

Disturbance severity: **0.48**

Disturbance severity (herb layer): **0.46**

Mowing frequency: **0.61**

Grazing pressure: **0.32**

Soil disturbance: **0.2**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#), [LA \(HER\) Salicetea herbaceae](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland**

Distribution

Continentality: **6**

Continentality amplitude: **6**