Festuca braun-blanquetii

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **8**
Moisture indicator value: **6**
Reaction indicator value: **4**
Nutrient indicator value: **1**

Disturbance indicator values

Disturbance frequency: **0.05**
Disturbance frequency (herb layer): **1.29**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.11**
Mowing frequency: **0.01**
Grazing pressure: **0.2**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CN (SAB) *Junipero-Pinetea sylvestris***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T37 Mediterranean montane Pinus sylvestris-Pinus nigra forest**

Broad habitat

Occurrence in broad habitats: **Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz