

# *Festuca heteropachys*

## Habitus and growth type

Plant height [m]: **0.13**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **7.3**

Moisture indicator value: **2.5**

Reaction indicator value: **6**

Nutrient indicator value: **1**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.49**

Disturbance frequency (herb layer): **1.63**

Disturbance severity: **0.53**

Disturbance severity (herb layer): **0.32**

Mowing frequency: **0.17**

Grazing pressure: **0.29**

Soil disturbance: **0.19**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

## Distribution

Continentality: **3**

Continentality amplitude: **1**