

# *Gentiana pneumonanthe*

## Habitus and growth type

Plant height [m]: **0.3**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **16.44**

## Flower

Flowering period: **June-October**

## Fruit, seed and dispersal

Seed mass [mg]: **0.06**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.9**

Temperature indicator value: **5.2**

Moisture indicator value: **7.4**

Nutrient indicator value: **1.8**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.56**

Disturbance frequency (herb layer): **1.85**

Disturbance severity: **0.38**

Disturbance severity (herb layer): **0.37**

Mowing frequency: **0.6**

Grazing pressure: **0.27**

Soil disturbance: **0.11**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CI \(NAR\) \*Nardetea strictae\*](#), [CM \(MOL\) \*Molinio-Arrhenatheretea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R37 Temperate and boreal moist or wet oligotrophic grassland](#), [S41 Wet heath](#)

Constant species of EUNIS habitats: [R37 Temperate and boreal moist or wet oligotrophic grassland](#), [S41 Wet heath](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality amplitude: 7